Oxidation-Reaction Reactions

- 1) What is oxidation?
- 2) What is reduction?
- What are the oxidation numbers in hydrogen peroxide, H_2O_2 ?
- 4) What are the oxidation numbers in sulfite, SO_3^{2-} ?
- 5) What determines whether a chemical reaction is a redox reaction?
- 6) (a) What effect does oxidation have on the oxidation number of a substance?
 - (b) What effect does reduction have on the oxidation number of a substance?
- 7) What is the difference between an oxidizing agent and a reducing agent?
- 8) Why are the Group I elements (alkali metals) very strong reducing agents?
- 9) Why are the Group VII elements (halogens) very strong oxidizing agents?
- 10) What is the relationship between the strength of an oxidizing agent and its corresponding reducing agent?
- 11) In the reaction, $O_2 + 4e^- \rightarrow 2O^{2-}$, is O_2 oxidized or reduced?
- 12) In the reaction, $2Na + Cl_2 \rightarrow 2NaCl$, which species is reduced?
- 13) In the reaction, $Mg + F_2 \rightarrow MgF_2$, which species is oxidized?
- In a redox reaction, MnO₄ is changed to Mn²⁺. How many electrons must be lost or gained by Mn?
- 15) In a redox reaction, MnO₄⁻ is changed to MnO₄²-. How many electrons must be lost or gained by Mn?
- In a balanced redox equation, how does the total number of reactant molecules compare with the total number of product molecules?
- 17) What is the strongest reducing agent among the elements?
- 18) What is the strongest oxidizing agent among the elements?

- 19) Balance the following redox reaction using the oxidation number method. $Al + Cu^{2+} \rightarrow Al^{3+} + Cu$
- 20) Balance the following redox reaction using the oxidation number method.

 copper + nitric acid → copper(II) nitrate + nitrogen monoxide + water

Solutions

- 1) Oxidation is a chemical reaction in which the atoms or ions of an element acquire a more positive oxidation state by losing electrons.
- 2) Reduction is a chemical reaction in which the atoms or ions of an element acquire a more negative oxidation state by gaining electrons.
- 3) +1 -1 H_2O_2
- 4) +4 -6 -2 SO₃²-
- 5) Only those reactions in which elements undergo a change in oxidation number (state).
- 6) (a) The oxidation number becomes more positive even though it may remain negative.
 - (b) The oxidation number becomes more negative even though it may remain positive.
- 7) The oxidizing agent is reduced or acquires a more negative oxidation state and the reducing agent is oxidized or acquires a more positive oxidation state.
- 8) The relatively large atomic radii of the alkali metals results in a weak force of attraction between the positive nuclei and the one valence electron.
- 9) The relatively small atomic radii of the halogens results in a strong force of attraction between the positive nuclei and the seven valence electrons.
- 10) The stronger an oxidizing agent, the weaker its corresponding reducing agent.

11)	O2 is reduced because its oxidation state becomes more negative by changing
	from 0 to -2.

- Cl₂ is reduced because its oxidation state becomes more negative by changing **12**) from 0 to -1.
- 13) Mg is oxidized because its oxidation state becomes more positive by changing from 0 to +2.

 $MnO_4^- \rightarrow Mn^{2+}$, therefore Mn must gain five electrons. **14**)

- **15**)
- **16**) As in all chemical reactions, there is no relationship that exists between the number of reactant and product molecules.
- **17**) Lithium.
- Fluorine. **18**)

19)
$$Al + Cu^{2+} \rightarrow Al^{3+} + Cu$$

$$0 +2 +3 0 \\ Al + Cu^{2+} \rightarrow Al^{3+} + Cu$$

oxidation:

$$Al \rightarrow Al^{3+} + 3e^{-}$$

reduction:

$$Cu^{2+} + 2e^{-} \rightarrow Cu$$

$$2Al \rightarrow 2Al^{3+} + 6e^{-}$$

$$3Cu^{2+} + 6e^{-} \rightarrow 3Cu$$

$$2Al + 3Cu^{2+} \Rightarrow 2Al^{3+} + 3Cu$$

$$2Al(s) + 3Cu^{2+}(aq) \Rightarrow 2Al^{3+}(aq) + 3Cu(s)$$

20) copper + nitric acid → copper(II) nitrate + nitrogen monoxide + water

$$Cu + HNO_3 \rightarrow Cu(NO_3)_2 + NO + H_2O$$

$$+5 -6$$
 $0 +1 -2 +2 +5 -2 +2 -2 +1 -2$
 $Cu + HNO_3 \rightarrow Cu(NO_3)_2 + NO + H_2O$

oxidation:

$$Cu \rightarrow Cu^{2+} + 2e^{-}$$

reduction:

$$N^{5+} + 3e^{-} \rightarrow N^{2+}$$

$$3Cu \rightarrow 3Cu^{2+} + 6e^{-}$$

$$2N^{5+} + 6e^{-} \rightarrow 2N^{2+}$$

 $3Cu + 2N^{5+} \rightarrow 3Cu^{2+} + 2N^{2+}$

$$3Cu + 2HNO_3 \rightarrow 3Cu(NO_3)_2 + 2NO + H_2O$$

$$3Cu + 8HNO_3 \rightarrow 3Cu(NO_3)_2 + 2NO + 4H_2O$$

 $3Cu(s) + 8HNO_3(aq) \rightarrow 3Cu(NO_3)_2(aq) + 2NO(g) + 4H_2O(l)$