

NEW PROOF OF THE SYMMEDIAN POINT TO BE THE CENTROID OF ITS PEDAL TRIANGLE, AND THE CONVERSE

Darij Grinberg

The following theorem is an important property of the symmedian point of a triangle (Fig. 1):




Fig. 1




Fig. 2

Theorem 1: Let L be the symmedian point of a triangle ABC . From L , drop the perpendiculars LX , LY , LZ on the sides BC , CA , AB , where X , Y , Z are the respective feet of the perpendicular. Then L is the centroid of the triangle XYZ .

We mention that in the customary terminology, the triangle XYZ is called the pedal triangle of L with respect to the triangle ABC ; but the triangle XYZ is also called **Lemoine**

pedal triangle of triangle ABC .

Theorem 1 can be paraphrased as follows: The symmedian point of a triangle is the centroid of the Lemoine pedal triangle.

Several proofs of Theorem 1 are known. In [2], p. 72-74, two proofs are presented. The proof given in [1] is a standard synthetic proof by constructing an auxiliary triangle.




Fig. 3

We will prove Theorem 1 with the help of another construction; in fact, we regard the intersection D of the tangents to the circumcircle of $\triangle ABC$ through the vertices B and C (Fig. 2). After [2], p. 60, the point D lies on the symmedian from the vertex A , i. e. on the symmedian AL . Thus, we have:

Lemma 2: The symmedian point L lies on the line AD .

Now we will prove the following lemma (Fig. 3):

Lemma 3: Drop perpendiculars DX' , DY' , DZ' from D to the sides BC , CA , AB . Then $DY'X'Z'$ is a parallelogram. [This theorem is interesting to have another signification: It means that an ex-symmedian point D is an ex-centroid (exmedian point) of its pedal triangle $X'Y'Z'$.]

Proof (Fig. 4): We denote the angles of triangle ABC by $\angle CAB = \alpha$, $\angle ABC = \beta$ and $\angle BCA = \gamma$. As chord-tangent angles, the angles $\angle CBD$ and $\angle BCD$ are both equal to the chordal angle of the chord BC , i. e. the angle α . From this, we have

$$\angle DBZ' = 180^\circ - \angle ABC - \angle CBD = 180^\circ - \beta - \alpha = \gamma.$$


Since $\angle DX'B = 90^\circ$ and $\angle DZ'B = 90^\circ$, the points X' and Z' lie on the circle having the segment DB as diameter, and consequently, $BX'DZ'$ is a cyclic quadrilateral, and as chordal angles $\angle DX'Z' = \angle DBZ'$. Thus,

$$\angle DX'Z' = \gamma. \quad (1)$$

Since the points X' and Y' lie on the circle having the segment DC as diameter (because $\angle DX'C = 90^\circ$ and $\angle DY'C = 90^\circ$), $CX'DY'$ is a cyclic quadrilateral, and this yields

By comparison with (1), we get $\triangle DX'Z' = \triangle X'DY'$, and from this, $X'Z' \parallel DY'$. Analogously, we can prove $X'Y' \parallel DZ'$, and thus, $DY'X'Z'$ is a parallelogram, qed.

Analogously, we can prove $X'Y' \parallel DZ'$, and thus, $DY'X'Z'$ is a parallelogram, qed.



Note that from the parallelogram $DY'X'Z'$, we get: The diagonal DX' bisects the diagonal $Y'Z'$. That means that the D -median of triangle $DY'Z'$ is DX' . Since DX' is orthogonal to BC , we have:

Now we will connect this with Theorem 1 (Fig. 5). Since L lies on AD and Y lies on AY' , and $LY \parallel DY'$ (because $LY \perp CA$ and $DY' \perp CA$), we have $AY : AY' = AL : AD$. Similarly, $AZ : AZ' = AL : AD$, and we conclude $AY : AY' = AZ : AZ'$. This yields $YZ \parallel Y'Z'$. Thus, the corresponding sides of triangles LYZ and $DY'Z'$ are parallel ($LY \parallel DY'$, $YZ \parallel Y'Z'$ and $ZL \parallel Z'L'$); therefore, also the L -median of triangle LYZ is parallel to the D -median of triangle $DY'Z'$. After Lemma 4, the latter one is orthogonal to BC ; thus, also the L -median of triangle LYZ is orthogonal to BC , i. e. the perpendicular from L to BC bisects the segment YZ . But this perpendicular is the line LX . Thus, LX bisects the segment YZ , i. e. in the triangle XYZ , LX is a median. Similarly, LY and LZ are the two other medians in triangle XYZ , and therefore, L is the centroid of triangle XYZ . This proves Theorem 1.




Fig. 5

This was my new proof. Also interesting is the observation that Theorem 1 possesses a valid converse theorem. We will now present a proof of *Theorem 1 together with the converse*:

Theorem 5 (Theorem 1 together with the converse): Let P be an arbitrary point in the plane of a triangle ABC , and let $\triangle XYZ$ be the pedal triangle of P with respect to triangle ABC . Then P is the centroid of triangle XYZ if and only if P is the symmedian point of triangle ABC .

In other words: There exists only one point which is the centroid of its pedal triangle, and this is the symmedian point.

The *proof* is similar to the trigonometric proof of Theorem 1 in [2], p. 72-73.

For first, we need the following lemma (proven in [2], p. 59):

Lemma 6: The distances of a point to the sides of a triangle ABC are in the ratios of these sides if and only if the point is the symmedian point of the triangle. In other words: For the distances $x = PX$, $y = PY$, $z = PZ$ of P to BC , CA , AB , the equation

$$x : y : z = a : b : c$$

holds if and only if P coincides with the symmedian point L of $\triangle ABC$.

Now we want to see when the point P is the centroid of its pedal triangle XYZ .

When does PY bisect the segment ZX ? Let D be the intersection of PY and ZX . Then, after the sine law in triangles PDZ and PDX , we have

$$\frac{ZD}{DX} = \frac{\sin \angle ZPD \cdot PZ : \sin \angle PDZ}{\sin \angle XPD \cdot PX : \sin \angle PDX} = \frac{\sin \angle ZPD}{\sin \angle XPD} \cdot \frac{PZ}{PX} : \frac{\sin \angle PDZ}{\sin \angle PDX}.$$

The angles $\angle PDZ$ and $\angle PDX$ sum up to 180° ; thus, their sines are equal:

$\sin \angle PDZ = \sin \angle PDX$, and we get

$$\frac{ZD}{DX} = \frac{\sin \angle ZPD}{\sin \angle XPD} \cdot \frac{PZ}{PX} = \frac{\sin \angle ZPD}{\sin \angle XPD} \cdot \frac{z}{x}. \quad (2)$$

For the angle $\angle ZPD$, we have $\angle ZPD = 180^\circ - \angle ZPY$; but we also have $\angle ZAY = 180^\circ - \angle ZPY$, since $AZPY$ is a cyclic quadrangle (as for $\angle AZP = 90^\circ$ and

$\angle AYP = 90^\circ$, the points Z and Y lie on the circle having segment AP as diameter). This yields $\angle ZPD = \angle ZAY$, or $\angle ZPD = \alpha$. Analogously, one finds $\angle XPD = \gamma$; with this, the equation (2) is simplified to

$$\frac{ZD}{DX} = \frac{\sin \alpha}{\sin \gamma} \cdot \frac{z}{x} = \frac{a}{c} \cdot \frac{z}{x} = \frac{z}{x} : \frac{c}{a}.$$

Therefore, $ZD = DX$ holds if and only if $z : x = c : a$. This means that P lies on the Y -median of triangle XYZ if and only if $z : x = c : a$. Analogously, P lies on the X -median of triangle XYZ if and only if $y : z = b : c$. Thus, P is the centroid of triangle XYZ (lies on two medians) if and only if $x : y : z = a : b : c$. After Lemma 6, the condition $x : y : z = a : b : c$ holds if and only if P is the symmedian point of $\triangle ABC$. Therefore, P is the centroid of the pedal triangle XYZ if and only if P is the symmedian point of $\triangle ABC$. This proves Theorem 5.




Fig. 6

References

- [1] Emil Donath: *Die merkwürdigen Punkte und Linien des ebenen Dreiecks*, Berlin 1976.
- [2] Ross Honsberger: *Episodes in Nineteenth and Twentieth Century Euclidean Geometry*, USA 1995.